

Symbol	“Think”	Value of L^\dagger	Formal Meaning [‡]
\circ	“ $<$ ”	$L = 0$	$f(n) \in O(g(n))$ but not $f(n) \in \Omega(g(n))$
\mathbf{O}	“ \leq ”	$L < \infty$	$\exists c, n_0[\forall n \geq n_0[f(n) \leq cg(n)]]$
$\mathbf{\Omega}$	“ \geq ”	$L > 0$	$\exists c, n_0[c \neq 0 \ \& \ \forall n \geq n_0[f(n) \geq cg(n)]]$
$\mathbf{\Theta}$	“ $=$ ”	$0 < L < \infty$	$f(n) \in O(g(n))$ and also $f(n) \in \Omega(g(n))$

[†] Value of $L = \lim_{n \rightarrow \infty} f(n)/g(n)$ when it exists.

[‡] Formal meaning of $f(n) = _ (g(n))$.

Figure 1: Greek Notation

Asymptotic Formulæ

$f(n) \in o(g(n))$ (old style $f(n) = o(g(n))$)

Said: $f(n)$ is little Oh of $g(n)$

Casual: f is *strictly* smaller (faster) than g .

Definition: $f(n) \in O(g(n))$ but not $f(n) \in \Omega(g(n))$

Calculus: $\lim_{n \rightarrow \infty} f(n)/g(n) = 0$ if the limit exists.

$f(n) \in O(g(n))$ (old style $f(n) = O(g(n))$)

Said: $f(n)$ is big Oh of $g(n)$

Casual: f is smaller (faster) than or equal to g .

Informal: f is eventually smaller than g (up to a constant).

Definition: $\exists c, n_0$ such that $\forall n \geq n_0 \quad f(n) \leq cg(n)$

Calculus: $\lim_{n \rightarrow \infty} f(n)/g(n) < \infty$ if the limit exists.

$f(n) \in \Omega(g(n))$ (old style $f(n) = \Omega(g(n))$)

Said: $f(n)$ is big Omega of $g(n)$

Casual: f is greater (slower) than or equal to g .

Informal: f is eventually greater than g (up to a constant).

Definition: $\exists c \neq 0, n_0$ such that $\forall n \geq n_0 \quad f(n) \geq cg(n)$

Calculus: $\lim_{n \rightarrow \infty} f(n)/g(n) > 0$ if the limit exists.

$f(n) \in \Theta(g(n))$ (old style $f(n) = \Theta(g(n))$)

Said: $f(n)$ is big Theta of $g(n)$

Casual: f is approximately the same as g .

Definition: $f(n) \in O(g(n))$ and also $f(n) \in \Omega(g(n))$

Calculus: $\lim_{n \rightarrow \infty} f(n)/g(n) \in (0, \infty)$ if the limit exists.

Usage the symbols O , o , Ω , and Θ are *not* function symbols. Formally $\Omega(h)$ is a class of functions. Used “old style” they should only be used “at the top level” on the right hand side of an equation. **Always use parentheses after these symbols.**

Generic $\Theta(f(n))$ formulæ.

- If $f(n) = c \cdot g(n)$ then $f(n) \in \Theta(g(n))$. *Constants don't matter.*
- If $f(n) = a_0 + a_1n + a_2n^2 + \dots + a_{k-1}n^{k-1} + a_kn^k$, where $a_k \neq 0$, then $f(n) \in \Theta(n^k)$. *Don't sweat the small stuff (polynomials).*
- If $g(n) \in o(f(n))$ then $f(n) + g(n) \in \Theta(f(n))$. *Don't sweat the small stuff (general).*
- $f(n) + g(n) \in \Theta(\max(f(n), g(n)))$. [This one comes up over and over in algorithm analysis.]
- If $f(n) \in \Theta(\log n)$ and $g(n) \in \Theta(n^\epsilon)$ (where $\epsilon > 0$) then $f(n) \in o(g(n))$. *Logarithms are smaller than polynomials.*
- If $f(n) \in \Theta(n^k)$ and $g(n) \in \Theta(c^n)$, where $c > 1$, then $f(n) \in o(g(n))$. *Polynomials are smaller than exponential functions.*

Logarithms.

- If $f(n) \rightarrow +\infty$ and $g(n) \rightarrow +\infty$ and $f(n) \in \Theta(g(n))$ then $\log f(n) \in \Theta(\log g(n))$. (*N.B. the converse does not hold!*) *logs preserve Θ .*
- [Charlie's Rule] If $f(n) \rightarrow +\infty$ and $g(n) \rightarrow +\infty$ and $\lim_{n \rightarrow \infty} \frac{\log f(n)}{\log g(n)} < 1$ then $f(n) \in o(g(n))$.¹ This rule helps a lot in dealing with odd functions like $n^{1/(\log \log n)}$.
- $\log_a n \in \Theta(\log_b n)$. *The base doesn't matter.*

Factorials.

Stirling's formula $n! = \sqrt{2\pi n} \left(\frac{n}{e}\right)^n \left(1 + \frac{1}{12n} + \frac{1}{288n^2} + g(n)\right)$

where $g(n) \in o(n^{-2})$.

Consequently $\log n! = \Theta(n \log n)$.

¹I know about this nice result from Charlie Obimbo, who presented it at WCCCE '03.