ASSIGNMENT 10 CHEMISTRY 200 Due: 4:30 pm Friday 28 November 2008

The Entropy Assignment

- 1. Consider the adiabatic irreversible compression of 9.66 moles of a ideal gas initially at a volume of 26.8 L and 369 K. This gas is compressed at constant pressure to half its initial volume.
 - (a) What are the final pressure and temperature?
 - (b) Calculate the entropy change associated with this change in state by each of the following combinations of reversible paths:
 - i. an isochore followed by an isobar
 - ii. an isotherm followed by an isochore
 - iii. an isotherm followed by an isobar
 - iv. an adiabat followed by an isochore
 - v. an adiabat followed by an isobar
 - vi. an abiabat followed by an isotherm
- 2. The molar heat capacity of gaseous hydrogen is given by $C_p/R = a + bT + cT^2$ where $a = 3.4958, b = -1.006 \times 10^{-4} \text{ K}^{-1}$, and $c = 2.419 \times 10^{-7} \text{ K}^{-2}$.
 - (a) What is the entropy change if the gas is heated at constant pressure from 300 K to 1500 K?
 - (b) What is the entropy change if the gas is heated at constant volume from 300 K to 1500 K?
 - (c) The heat capacity of a gas can be estimated by taking into account the internal motions of the molecule and assuming that there are no intermolecular forces. Each vibrational degree of freedom contributes R to the molar heat capacity while each rotational and each translational degree of freedom contributes R/2. On this basis, recalculate the entropy changes, assuming no temperature dependence. Discuss whether the vibrational motion of the molecule affects the heat capacity.
- 3. The difference between heat capacity at constant pressure and at constant volume is given by:

$$C_p - C_v = T\left(\frac{\partial V}{\partial T}\right)_p \left(\frac{\partial p}{\partial T}\right)_v$$

- (a) Evaluate this expression for a perfect gas.
- (b) Evaluate this expression for a hard sphere gas.
- (c) Evaluate this expression for a gas with a Berthelot equation of state.

$$P = \frac{nRT}{V - nb} - \frac{n^2a}{TV^2}$$

HINT: Use the cyclic rule

$$-1 = \left(\frac{\partial V}{\partial T}\right)_p \left(\frac{\partial T}{\partial p}\right)_V \left(\frac{\partial p}{\partial V}\right)_T$$

as necessary.

- 4. One mole of perfect gas doubles in volume at a constant temperature. What volume change is required for 2.00 mole of an ideal gas to exhibit the same increase in entropy? For 0.500 mol?
- 5. Calculate the absolute entropy of one mole of S gas at 917 K. At 95.4°C, the enthalpy of the transition from rhombic to monoclinic form is 0.38 kJ mol⁻¹. At 119°C, monoclinic S melts with an enthalpy of fusion of 1.23 kJ mol⁻¹. The boiling point is 444.7°C. The heat capacity of rhombic S between 15 and 370 K is:

$$C_p/R = 0.08 + 2.17 \times 10^{-2}T - 6.2 \times 10^{-5}T^2 + 6.9 \times 10^{-8}T^3$$

Below 15 K, the Debye expression is used:

$$\frac{C_v}{R} = \frac{233.8}{\Theta_D^3} T^3$$

where the Debye temperature, Θ_D for rhombic S is 150.4 K. The heat capacity of monoclinic S is:

$$C_p/R = 1.96 \times 10^{-2}T - 4.8 \times 10^{-5}T^2 + 4.7 \times 10^{-8}T^3$$

The heat capacity of liquid S is:

$$C_p/R = 7.85$$

Use Trouton's rule:

$$\frac{\Delta H_{vap}}{T_{vap}} = 88 \text{ J K}^{-1} \text{ mol}^{-1}$$

to estimate the enthalpy of vaporization.

In the gas phase, assume that atomic S behaves ideally.