


GSA TODAY

VOL. 19, No. 8

A PUBLICATION OF THE GEOLOGICAL SOCIETY OF AMERICA

AUGUST 2009

The Klondike goldfields and Pleistocene environments of Beringia



Inside:

- ▲ **GSA-USGS Congressional Science Fellow Named for 2009–2010**, p. 13
- ▲ **Field Forum Scheduled:** Significance of along-strike variations for the 3-D architecture of orogens: The Hellenides and Anatolides in the eastern Mediterranean, p. 14
- ▲ **Groundwork:** Mineral resource geology in academia: An impending crisis?, p. 27

rockware full page color ad here

4 **The Klondike goldfields and Pleistocene environments of Beringia**

Duane G. Froese, Grant D. Zazula, John A. Westgate, Shari J. Preece, Paul T. Sanborn, Alberto V. Reyes, and Nicholas J.G. Pearce



Cover: Following the footsteps of stampeders to the Klondike creeks, scientists from across Canada and abroad were lured to the area by reports of spectacular ice-age fossils from the permafrost, including this mammoth skull (not mastodon as written) recovered from Quartz Creek following the Gold Rush. Photo courtesy Dawson City Museum Archives. See "The Klondike goldfields and Pleistocene environments of Beringia" by D.G. Froese et al., p. 4–10.

12 **Letter**

13 **GSA-USGS Congressional Science Fellow Named for 2009–2010**

14 **Field Forum Scheduled:** Significance of along-strike variations for the 3-D architecture of orogens: The Hellenides and Anatolides in the eastern Mediterranean

17 **2009 GSA Annual Meeting Luncheon Keynote Lectures**

20 **2009 GSA Mentor Program Highlights**

22 **GSA Foundation Update**

23 **GSA Awards 2008 Intel International Science and Engineering Fair Prize**

24 **Calendar of Upcoming GSA Deadlines & Events**

24 **Notice of GSA Council Meeting**

24 **Earth Science Literacy Initiative**

25 **Classified Advertising**

27 **Groundwork:** Mineral resource geology in academia: An impending crisis?

29 **Geologic Past—**“The Role of Minerals in the Present International Situation”

30 **Journal Highlights**

GSA TODAY publishes news and information for more than 22,000 GSA members and subscribing libraries. *GSA TODAY* (ISSN 1052-5173 USPS 0456-530) is published 11 times per year, monthly, with a combined April/May issue, by The Geological Society of America®, Inc., with offices at 3300 Penrose Place, Boulder, Colorado. Mailing address: P.O. Box 9140, Boulder, CO 80301-9140, USA. Periodicals postage paid at Boulder, Colorado, and at additional mailing offices. Postmaster: Send address changes to *GSA Today*, GSA Sales and Service, P.O. Box 9140, Boulder, CO 80301-9140, USA. GSA provides this and other forums for the presentation of diverse opinions and positions by scientists worldwide, regardless of their race, citizenship, gender, religion, or political viewpoint. Opinions presented in this publication do not reflect official positions of the Society.

Copyright © 2009, The Geological Society of America (GSA). All rights reserved. Copyright not claimed on content prepared wholly by U.S. government employees within the scope of their employment. Individual scientists are hereby granted permission, without fees or further requests to GSA, to use a single figure, a single table, and/or a brief paragraph of text in other subsequent works and to make unlimited photocopies of items in this journal for noncommercial use in classrooms to further education and science. For any other use, contact Permissions, GSA, P.O. Box 9140, Boulder, CO 80301-9140, USA, Fax +1-303-357-1073, editing@geosociety.org.

SUBSCRIPTIONS for the 2009 calendar year: Society Members: *GSA Today* is provided as part of membership dues. Contact GSA Sales and Service at +1-888-443-4472, +1-303-357-1000, option 3, or gsaservice@geosociety.org for membership information. Nonmembers & Institutions: Free with paid subscription to both *GSA Bulletin* and *Geology*, otherwise US\$60. Contact AIP Customer Service, subs@aip.org. Claims: For nonreceipt or for damaged copies, members contact GSA Sales and Service; all others contact AIP Customer Service, subs@aip.org. Claims are honored for one year; please allow sufficient delivery time for overseas copies, up to six months.

GSA TODAY STAFF:

Executive Director and Publisher: John W. Hess

Science Editors: Stephen T. Johnston, University of Victoria, School of Earth & Ocean Sciences, Victoria, British Columbia V8W 3P6, Canada, stj@uvic.ca; and David E. Fastovsky, University of Rhode Island, Department of Geosciences, Woodward Hall, Rm. 317, Kingston, Rhode Island 02881, USA, defastov@uri.edu.

Managing Editor: Kristen E. Asmus, kasmus@geosociety.org

Graphics Production: Margo Y. Sajban

ADVERTISING:

Classifieds & Display: Ann Crawford, +1-800-472-1988, ext. 1053, +1-303-357-1053, Fax +1-303-357-1070; acrawford@geosociety.org

GSA ONLINE: www.geosociety.org

Printed in the USA using pure soy inks.

The Klondike goldfields and Pleistocene environments of Beringia

Duane G. Froese, Dept. of Earth and Atmospheric Sciences, Univ. of Alberta, Edmonton, Alberta T6G 2R3; **Grant D. Zazula**, Yukon Palaeontology Program, Whitehorse, Yukon Territory Y1A 2C6; **John A. Westgate**, **Shari J. Preece**, Dept. of Geology, Univ. of Toronto, Toronto, Ontario M5S 3B1; **Paul T. Sanborn**, Ecosystem Science and Management Program, Univ. of Northern British Columbia, Prince George, British Columbia V2N 4Z9; **Alberto V. Reyes**, Dept. of Earth and Atmospheric Sciences, Univ. of Alberta, Edmonton, Alberta T6G 2R3; and **Nicholas J.G. Pearce**, Inst. of Geography and Earth Science, University of Wales, Aberystwyth SY23 3DB, UK

ABSTRACT

The Klondike goldfields of Yukon, Canada, contain a key record of Pleistocene Beringia, the region of Yukon, Alaska, and Siberia that remained largely unglaciated during the late Cenozoic. A concentration of mining exposures, with relict permafrost that is locally more than 700,000 years old, provides exceptional preservation of paleoenvironmental archives and a new perspective on the nature of paleoenvironments during the Pleistocene. A critical feature is the stratigraphic association of distal tephra beds with these paleoenvironmental ar-

chives, which facilitates their regional correlation and, in many cases, provides independent ages for the paleoenvironmental assemblages. Paleoenvironmental analyses of fossil arctic ground-squirrel middens and buried vegetation indicate the presence of cryoxerophilous ("steppe-tundra") vegetation growing on well-drained substrates with deep active layers (seasonal thaw depths) during cold intervals of the Pleistocene. Studies of full-glacial paleosols and cryostratigraphic relations of associated ground ice indicate the importance of active loess deposition and surface vegetation cover in maintaining the functionally distinct mammoth-steppe biome, which supported grazing mega-fauna populations, including mammoth, horse, and bison.

INTRODUCTION

Swedish biogeographer Eric Hultén introduced the concept of Beringia to explain the distribution of arctic and boreal plants around the Bering Strait. He proposed that a continuous Holarctic refugium beyond the continental ice sheets of North America existed during the Quaternary (Hultén, 1937). Hultén originally considered Beringia as the region of the continental shelf that was exposed when lowered sea level connected eastern Asia with North America, but we now consider it more




Figure 1. Eastern Beringia during the last glacial maximum with eustatic sea level lowering of 120 m. The region was largely unglaciated with the exception of local uplands that supported alpine glaciers (glacier limits from Ehlers and Gibbard, 2004).

broadly to include the unglaciated landmass from the Kolyma River in Siberia to Yukon, Canada. This area extends some 3000 km across, including the Klondike region toward the eastern edge of that boundary (Fig. 1).

Beringia represents the largest contiguous Arctic expanse to remain unglaciated during the Pliocene and Pleistocene and preserves an exceptional sedimentary archive spanning the past several million years. Relict permafrost (Kotler and Burn, 2000; Froese et al., 2008) in Beringia has preserved a diversity of exceptional paleoenvironmental archives, including mammals (Guthrie, 1990), paleobotanical remains (Goetcheus and Birks, 2001; Zazula et al., 2003), and ancient DNA (Shapiro et al., 2004). Critical to stratigraphic integrity and to understanding the paleoenvironmental significance of these archives is the presence of widespread ($>10^6$ km²) and numerous distal volcanic ash layers or tephra beds (Westgate et al., 2001). These tephra beds are datable by a variety of methods, including glass fission-track (Westgate et al., 2001) and associated radiocarbon ages (Froese et al., 2002), but they also provide correlative timelines between sites based on their unique geochemistry. The presence of these tephra beds within the perennially frozen late Cenozoic sediments of eastern Beringia is unique to this area of the northern hemisphere. Similar permafrost-preserved records are found in Siberia (e.g., Sher et al., 2005), but the lack of readily datable materials there, namely a rich tephrostratigraphy, is a challenge to developing chronologies beyond the range of radiocarbon dating (ca. 50,000 years).

Substantial progress has been made since Hultén's time in documenting the biogeographic significance of Beringia (Hopkins et al., 1982; Brigham-Grette, 2001), including its role as an evolutionary center (Sher, 1997) and as the crossroads for faunal exchanges between Asia and North America (e.g., Sher, 1999; Repenning and Brouwers, 1992; Shapiro et al., 2004). However, considerable debate has focused on how Beringia could support a diverse grazing megafauna under Pleistocene glacial climates (Guthrie, 1990). Discussion is based largely on interpretations of lacustrine paleoecological records that differ from those derived from river bluff exposures and, especially, exposures in the Fairbanks placer mining region (Hopkins et al., 1982; Guthrie, 1990). Until recently, however, the Klondike region has received comparatively little attention beyond the analysis of late Pleistocene vertebrate fossils (Harington, 2003). Here, we highlight recent study of the tephrostratigraphy and associated paleoenvironmental archives in the Klondike region, which has led to new understanding of late Pleistocene environments of Beringia.

REGIONAL SETTING

The interior of Yukon and Alaska has a strongly continental climate due to the pronounced rain shadow of the Coast and St. Elias Mountains of western Canada and the Alaska Range and Wrangell mountains of southern Alaska (Fig. 1). This aridity was likely established by the Pliocene (White et al., 1997; Froese et al., 2000), such that during Plio-Pleistocene glacial intervals, the interior of Yukon and Alaska was cold enough to support ice sheets but too dry for extensive glaciation. The Klondike region lies at the eastern edge of this unglaciated area, within 150 km of the last glacial maximum Cordilleran Ice Sheet (Fig. 1).



Figure 2. *Bison priscus* skull recovered from sediments associated with the late Pleistocene Dawson tephra on a tributary to Dominion Creek. Pleistocene fossils are still actively recovered from perennially frozen placer mining exposures in the Klondike, particularly in narrow valleys and near hillslopes where loessal “muck” deposits have aggraded.

Since the discovery of placer gold in the Klondike in 1896 and the subsequent gold rush, mining has produced tremendous exposures of surficial sediments within the Klondike goldfields along with the recognition of abundant Pleistocene fossil bones (e.g., Harington and Clulow, 1973). Prior to the gold rush, G.M. Dawson and R.G. McConnell of the Geological Survey of Canada (Dawson, 1894) had collected fossils from the area, and the Muséum d'histoire naturelle de Paris, the U.S. Biological Survey, and the American Museum of Natural History sent researchers to collect ice-age fossils in the early 1900s. Perhaps the most noteworthy of all paleontologists to have worked in the Klondike is C.R. Harington, who made sizable vertebrate collections for the Canadian Museum of Nature during the 1960s–1990s. Hundreds to thousands of fossils are still produced every year from placer gold mining and provide an invaluable research resource (Fig. 2).

Nearly all drainages of King Solomon Dome (Fig. 3) have produced gold, with total production estimated at ~15,000,000 ounces; active mining produces >50,000 ounces annually, largely from small, independently owned mines. Development of radiating drainage from King Solomon Dome (Fig. 3) during the Cenozoic released gold from bedrock sources, and, coupled with slow rates of uplift, produced prominent terraces in major valleys dating to the Pliocene and early Pleistocene (McConnell, 1907; Froese et al., 2000). In the valley bottoms, local creek gravels are associated with ice-rich loess, or “muck” deposits (Fig. 4).

TEPHROSTRATIGRAPHIC FRAMEWORK

The presence of numerous distal silicic tephra beds has been instrumental for the development and interpretation of the late Cenozoic sedimentary and paleoenvironmental record in the Klondike region (Preece et al., 2000; Westgate et al., 2001).




Figure 3. Klondike area map with locations of late Pleistocene Dawson tephra (after Froese et al., 2006)



Figure 4. Ice-rich loessal deposits, or “muck,” of the Klondike goldfields. Aggradation of loess with permafrost has led to exceptional preservation of paleoenvironmental archives in the Klondike area. Paleoenvironmental reconstruction from these muck deposits, sometimes called “Pleistocene in a blender,” was largely avoided because of their complexity. Detailed tephrostratigraphy has led to new understanding of their significance for reconstructing Pleistocene environments of eastern Beringia.




Figure 5. (A) Bivariate plot of $\text{SiO}_2\text{-K}_2\text{O}$ illustrating differences in geochemistry for some Klondike area tephra beds. (B) Rare earth element profiles for tephra beds shown in A. Data are normalized to chondrites using the values of Sun and McDonough (1989). Dawson, Old Crow, and Dominion Creek tephra beds have Type I characteristics, while Sheep Creek-K (Sheep Ck.-K) and White River Ash—east lobe (White R. E) have Type II characteristics. Gold Run tephra has a mixture of Type I and II characteristics, with glass trace-element compositions similar to Type I and mineralogy similar to Type II (data sources and analytical methods: Preece et al., 2000; Westgate et al., 2001, 2008; Pearce et al., 2004; this paper).

The glass morphology, mineral content, and geochemistry of each tephra bed help to reveal its provenance and suggest two broad volcanic source areas (Fig. 5; Preece et al., 2000). Tephra beds from the Aleutian arc–Alaska Peninsula, or Type I beds, have few crystals, mainly bubble-wall glass shards, abundant pyroxene, and rare earth element (REE) profiles with a well-developed negative Eu anomaly (Fig. 5). In contrast, Type II beds, derived from the Wrangell Volcanic Field (and Hayes volcano), have abundant crystals and glass that is mainly in the form of highly inflated pumice; hornblende is abundant, and REE profiles are steep, with a weakly developed Eu anomaly (Fig. 5). Physical and chemical properties, together with stratigraphic and paleoecological context, allow identification of tephra beds and their correlation between sites (Fig. 5). At least seven Type I beds are known in the Klondike, including Dawson (25.3 ^{14}C ka B.P.), Dominion Creek (82 \pm 9 ka), and Old Crow (131 \pm 11 ka); the 26 known Type II beds include White River Ash (1.2 and 1.7 ka) and Sheep Creek-K (ca. 80 ka). Six “other” beds are also present in the Klondike; these are either too mafic for classification or, like the Gold Run tephra (740 \pm 60 ka), have characteristics of both Type I and II tephra beds.

The most commonly observed bed is the late Pleistocene Dawson tephra, one of the largest Quaternary eruptions in eastern Beringia, likely exceeding 50 km^3 (Fig. 6; Froese et al., 2002, 2006; Mangan et al., 2003). Dawson tephra has been identified at more than 20 sites in the area (Fig. 3), where it typically occurs as a 30–80-cm-thick bed in “muck” deposits (Fig. 7). Bracketing radiocarbon ages on plant macrofossils provide a mean age of ca. 25,300 ^{14}C yr B.P. and a calendar year estimate of \sim 30,000 yr B.P. (Froese et al., 2006; Demuro et al., 2008); Dawson tephra marks the onset of glacial conditions of Marine Isotope Stage (MIS) 2 in central Yukon (Zazula et al., 2006). Other beds provide similar time markers in the Klondike: Old Crow tephra for late MIS 6 (131 \pm 11 ka; Péwé et al. 2009), directly below the last interglacial (MIS 5e) thaw unconformity; Sheep Creek-C tephra (ca. 90 ka) for late MIS 5 interglacial conditions; and Sheep Creek-K tephra (ca. 80 ka) for the MIS 5–4 transition (Westgate et al., 2008). Collectively, these and other beds provide key marker horizons for discrete timeslices between sites in the Klondike, allowing integration of diverse paleoenvironmental archives from relict permafrost, plant and insect macrofossils, pollen, and vertebrate remains.




Figure 6. Dawson and Old Crow tephra distribution (after Froese et al., 2002).



Figure 8. Ice wedge cross-cutting late MIS 5 forest bed at head level of person in photo along Dominion Creek. Sheep Creek tephra-K (ca. 80 ka) is present about midway through the silts (arrow), marking the late MIS 5–4 transition in central Yukon.



Figure 7. Late Pleistocene Dawson tephra (25,300 ^{14}C yr B.P.) overlying syngenetic ice wedge at Quartz Creek. Depression of tephra into the wedge top (upper right) marks the former depression of the polygonal ground network, indicating the ice wedge was active when the tephra was deposited. Arrows mark arctic ground squirrel middens; tephra thickness is ~40 cm.



Figure 9. Graminoid-rich paleosol with roots overlying ice wedge marking the paleo-active layer when the soil was formed. The paleosol is cross-cut by secondary ice wedge growth (vertical arrow at left) and includes beds of Dominion Creek tephra (82 ± 9 ka; horizontal arrow), marking early MIS 4 glacial conditions. Ice axe: 80 cm.

“MUCK”: BERINGIA’S ICE AGE FREEZER

Muck deposits in the Klondike are part of a broader complex of silts that blanket much of Beringia and are usually considered to be loess and locally re-transported loess from creeks and river valleys that aggraded with permafrost (Péwé, 1975; Fraser and Burn, 1997; Muhs et al., 2003; Fig. 8). In the discontinuous permafrost zone (50%–90% frozen ground), which includes the Klondike and Fairbanks regions, these ice-rich loessal deposits are found on north- and east-facing slopes and within narrow valleys along hillslopes. These frozen deposits have high organic carbon content (Fraser and Burn, 1997; Sanborn et al., 2006), reflecting aggradation of the permafrost table with silt deposition, and may reach tens of meters in thickness. These combined processes buried soils before their associated root detritus and other plant material could

decompose (Fig. 9), preserving Pleistocene organic remains in permafrost at some sites for >700,000 years (Froese et al., 2008). Similar Pleistocene deposits are present in Siberia, where they are termed Yedoma, though it is not well-accepted that these are, *sensu stricto*, eolian deposits (Sher, 1997; Schirmermeister et al., 2008).

The influence of surface vegetation cover on permafrost reveals important functional differences between the reconstructed Pleistocene glacial steppe-tundra environment (Zazula et al., 2003) and the modern boreal forest environment of interior Yukon and Alaska. Most sites with mucks present are north- and east-facing or in narrow valleys with black spruce (*Picea mariana*) forests and are characterized by thick covers of moss and partially humified vegetation litter. The thermal properties of this groundcover promote deep winter cooling

and insulate the ground from summer heat, resulting in poorly drained substrates with permafrost and shallow active layer depths. Recovery of MIS 2 and 4 arctic ground-squirrel middens and paleosols from muck deposits at these sites reveals that substrates were better drained during Pleistocene glacial intervals than they are at present (Zazula et al., 2005; Sanborn et al., 2006). In fact, arctic ground squirrels are absent from the Klondike region today, suggesting important expansion of their ranges during Pleistocene glacial intervals (Zazula et al., 2005). Present-day ground squirrels, in southern Yukon and the north slope of Alaska, require well-drained soils with active layer depths of ~1 m for burrowing and successful hibernation. The translocation of paleosol A-horizon material in the paleo-active layer and truncation of underlying ice bodies provide additional evidence for deeper active layers (Sanborn et al., 2006; Fig. 9). Thus, despite summer air temperature depression of several degrees Celsius during the glacial intervals (e.g., Elias, 2001), soils were better drained with deeper active layers due to the presence of graminoid vegetation cover, which lacked the insulating properties of modern soils in the region. Well-drained soils with deeper active layers and additions of soil nutrients from loess deposition would have enabled greater nutrient turnover, essential for a herbaceous steppe-tundra habitat that supported herbivores, such as woolly mammoths and horses (Laxton et al., 1996).

QUATERNARY MAMMAL FOSSILS

The Klondike is one of North America's most productive localities for the recovery of late Pleistocene mammal fossils (Harington, 2003). Most Klondike faunas are dominated by the "big-three" of Beringia—steppe bison (*Bison priscus*), woolly mammoth (Fig. 2; *Mammuthus primigenius*), and Yukon horse (*Equus lambei*). Fossils of less common species are recovered occasionally, including the western camel (*Camelops betherius*), American mastodon (*Mammuth americanum*), American lion (*Panthera leo atrox*), short-faced bear (*Arctodus simus*), and helmeted muskox (*Bootherium bombifrons*). Mummified or freeze-dried partial carcasses recovered from the Klondike highlight the role of permafrost in the preservation of the late Pleistocene paleontological record. Impressive mummified carcasses include black-footed ferret (*Mustela nigripes*) and Yukon horse (*Equus lambei*), whose stomach contents have provided important dietary information (Harington, 2007). The exceptional preservation of Klondike vertebrate bones has led to recent ancient biomolecule studies using mitochondrial DNA sequencing and radiocarbon dating to establish phylogenetic histories for bison (Shapiro et al., 2004), horse (Weinstock et al., 2005), and mammoth (Debruyne et al., 2008).

PALEOENVIRONMENTAL ARCHIVES

Questions concerning the nature of terrestrial ecosystems in Beringia have been a major research focus for Quaternary paleoecologists for decades (Hopkins et al. 1982; Guthrie, 1990; Birks and Birks, 2000). Although the Klondike region has been well known for Pleistocene mammal fossils for the past century, there has been little systematic paleoecological research in the region until the last decade. The abundance of Pleistocene vertebrate faunas and well-constrained stratigraphic records makes the Klondike a valuable region for resolving

questions concerning the relations between mammals, glacial vegetation, and Pleistocene climates.

Recent paleoecological work in the Klondike has focused on detailed analysis of fossil middens (nests, seed caches, and burrows) of arctic ground squirrels (*Spermophilus parryii*) (Zazula et al., 2003, 2007). In the Klondike, over 100 middens have been recovered and analyzed systematically in association with the Sheep Creek-K-Dominion Creek tephtras (ca. 80 ka) and Dawson tephra (ca. 25.3 ¹⁴C ka), providing paleoenvironmental records for MIS 4 and early MIS 2, respectively. Plant macrofossils (seeds, fruits, leaves) from the middens are dominated by grasses, dryland sedges, sage, and a wide variety of flowering forbs. Together, these plants formed an open, grass- and forb-rich steppe-tundra community that thrived on the well-drained, deeply thawed loessal soils in the Klondike during Pleistocene cold intervals.

CONCLUSIONS

The Klondike goldfields provide an exceptional record of Pleistocene Beringia. The development of a robust tephrostratigraphic and chronologic framework for the perennially frozen deposits has facilitated integration of paleoenvironmental archives from vertebrate remains, paleobotanical, paleosol, and cryostratigraphic observations. This mammoth-steppe environment was characterized by graminoid and forb-rich vegetation with better-drained loessal substrates and deeper active layers despite summer temperature depressions. Collectively, these records support the notion that functional differences between the cryoxeric steppe-tundra and the modern boreal environment provides a means to explain the existence of a rich grazing fauna during Pleistocene glacial intervals.

ACKNOWLEDGMENTS

We are indebted to the Klondike Placer Miners for unfettered access and generous assistance over the last 15 years of fieldwork in the Klondike area. Additional support for this research has been provided by the Natural Science and Engineering Research Council of Canada, Alberta Ingenuity, Geological Survey of Canada, Yukon Geological Survey, and Yukon Heritage. We also acknowledge the benefits of collaborative research and discussions with many colleagues on the Klondike record. Thanks also to former *GSA Today* Editor, Gerry Ross, for his initial suggestion of this paper, and Stephen Johnston for his patience in its receipt.

REFERENCES CITED

- Birks, H.H., and Birks, H.J.B., 2000, Future uses of pollen analysis must include plant macrofossils: *Journal of Biogeography*, v. 27, p. 31–35, doi: 10.1046/j.1365-2699.2000.00375.x.
- Brigham-Grette, J., 2001, New perspectives on Beringian Quaternary paleogeography, stratigraphy, and glacial history: *Quaternary Science Reviews*, v. 20, p. 15–24, doi: 10.1016/S0277-3791(00)00134-7.
- Dawson, G.M., 1894, Notes on the occurrence of mammoth remains in the Yukon district of Canada and in Alaska: *Quarterly Journal of the Geological Society of London*, v. 50, p. 1–9.
- Debruyne, R., Chu, G., King, C.E., Bos, K., Kuch, M., Schwarz, C., Szipak, P., Gröcke, D.R., Matheus, P., Zazula, G., Guthrie, D., Froese, D., Buigues, B., de Marliave, C., Flemming, C., Poinar, D., Fisher, D., Southon, J., Tikhonov, A.N., MacPhee, R.D.E., and Poinar, H.N., 2008, Out of America: Ancient DNA evidence for a New World origin of Late Quaternary woolly mammoths: *Current Biology*, v. 18, p. 1320–1326.
- Demuro, M., Roberts, R.G., Froese, D.G., Arnold, L.J., Brock, F., and Ramsey, C.B., 2008, Optically stimulated luminescence dating of

- single and multiple grains of quartz from perennially frozen loess in western Yukon Territory, Canada: Comparison with radiocarbon chronologies for the late Pleistocene Dawson tephra: *Quaternary Geochronology*, v. 3, p. 346–364.
- Ehlers, J., and Gibbard, P.L., 2004, Quaternary glaciations—Extent and chronology, Part II: North America. *Developments in Quaternary Science*: Amsterdam, Elsevier, 440 p.
- Elias, S.A., 2001, Mutual climate range reconstructions of seasonal temperatures based on Late Pleistocene fossil beetle assemblages in Eastern Beringia: *Quaternary Science Reviews*, v. 20, p. 77–91, doi: 10.1016/S0277-3791(00)00130-X.
- Fraser, T.A., and Burn, C.R., 1997, On the nature and origin of “muck” deposits in the Klondike area, Yukon Territory: *Canadian Journal of Earth Sciences*, v. 34, p. 1333–1344.
- Froese, D.G., Barendregt, R.W., Enkin, R.J., and Baker, J., 2000, Paleomagnetic evidence for multiple Late Pliocene–Early Pleistocene glaciations in the Klondike area, Yukon Territory: *Canadian Journal of Earth Sciences*, v. 37, p. 863–877, doi: 10.1139/cjes-37-6-863.
- Froese, D.G., Westgate, J.A., Preece, S.J., and Storer, J., 2002, Age and significance of the late Pleistocene Dawson tephra in eastern Beringia: *Quaternary Science Reviews*, v. 21, p. 2137–2142, doi: 10.1016/S0277-3791(02)00038-0.
- Froese, D.G., Zazula, G.D., and Reyes, A.V., 2006, Seasonality of the late Pleistocene Dawson tephra and exceptional preservation of a buried riparian surface in central Yukon Territory, Canada: *Quaternary Science Reviews*, v. 25, p. 1542–1551, doi: 10.1016/j.quascirev.2006.01.028.
- Froese, D.G., Westgate, J.A., Reyes, A.V., Enkin, R.J., and Preece, S.J., 2008, Ancient permafrost and a future, warmer arctic: *Science*, v. 321, p. 1648, doi: 10.1126/science.1157525.
- Goetcheus, V.G., and Birks, H.H., 2001, Full-glacial upland tundra vegetation preserved under tephra in the Beringia National Park, Seward Peninsula, Alaska: *Quaternary Science Reviews*, v. 20, p. 135–147, doi: 10.1016/S0277-3791(00)00127-X.
- Guthrie, R.D., 1990, *Frozen fauna of the mammoth steppe: the story of Blue Babe*: Chicago, University of Chicago Press, 323 p.
- Harington, C.R., 2007, Late Pleistocene mummified mammals, in Elias, S.A., ed., *Encyclopedia of Quaternary Science*: Amsterdam, Elsevier, p. 3197–3202.
- Harington, C.R., 2003, *Annotated bibliography of Quaternary vertebrates of northern North America—with radiocarbon dates*: Toronto, University of Toronto Press, 539 p.
- Harington, C.R., and Clulow, F.V., 1973, Pleistocene mammals from Gold Run Creek, Yukon Territory: *Canadian Journal of Earth Sciences*, v. 10, p. 697–759.
- Hopkins, D.M., Matthews, J.V., Jr., Schweger, C.E., and Young, S.B., eds., 1982, *Paleoecology of Beringia*: New York, Academic Press, 489 p.
- Hultén, E., 1937, *Outline of the history of arctic and boreal biota during the Quaternary period*: Stockholm, Thule, 168 p.
- Kotler, E., and Burn, C.R., 2000, Cryostratigraphy of the Klondike “muck” deposits, west-central Yukon Territory: *Canadian Journal of Earth Sciences*, v. 37, p. 849–861.
- Laxton, N.F., Burn, C.R., and Smith, C.A.S., 1996, Productivity of loessal grasslands in the Kluane Lake region, Yukon Territory, and the Beringian “production paradox”: *Arctic*, v. 49, p. 129–140.
- Mangan, M.T., Waythomas, C.F., Miller, T.P., and Trusdell, F.A., 2003, Emons Lake Volcanic Center, Alaska Peninsula: source of the Late Wisconsin Dawson tephra, Yukon Territory, Canada: *Canadian Journal of Earth Sciences*, v. 40, p. 925–936, doi: 10.1139/e03-026.
- Muhs, D.R., Ager, T.A., Arthur Bettis, E., III, McGeehin, J., Been, J.M., Béget, J.E., Pavich, M.J., Stafford, J., Thomas, W., and Stevens, D.A.S.P., 2003, Stratigraphy and palaeoclimatic significance of Late Quaternary loess-palaeosol sequences of the Last Interglacial-Glacial cycle in central Alaska: *Quaternary Science Reviews*, v. 22, p. 1947–1986, doi: 10.1016/S0277-3791(03)00167-7.
- McConnell, R.G., 1907, Report on gold values in the Klondike high level gravels: Geological Survey of Canada Publication 979, 34 p.
- Pearce, N.J.G., Westgate, J.A., Perkins, W.T., and Preece, S.J., 2004, The application of ICP-MS methods to tephrochronological problems: *Applied Geochemistry*, v. 19, p. 289–322, doi: 10.1016/S0883-2927(03)00153-7.
- Péwé, T.L., 1975, *Quaternary Geology of Alaska*: U.S. Geological Survey Professional Paper 835, 143 p.
- Péwé, T.L., Westgate, J.A., Preece, S.J., Brown, P.M., and Leavitt, S.W., 2009, Late Pliocene Dawson Cut Forest Bed and new tephrochronological findings in the Gold Hill Loess, east-central Alaska: *Geological Society of America Bulletin*, v. 121, p. 294–320.
- Preece, S.J., Westgate, J.A., Alloway, B.V., and Milner, M.W., 2000, Characterization, identity, distribution, and source of late Cenozoic tephra beds in the Klondike District of the Yukon, Canada: *Canadian Journal of Earth Sciences*, v. 37, p. 983–996, doi: 10.1139/cjes-37-7-983.
- Repenning, C.A., and Brouwers, E.M., 1992, Late Pliocene-early Pleistocene ecologic changes in the Arctic Ocean Borderland: *U.S. Geological Survey Bulletin* 2036, 37 p.
- Sanborn, P.T., Smith, C.A.S., Froese, D.G., Zazula, G.D., and Westgate, J.A., 2006, Full-glacial paleosols in perennially frozen loess sequences, Klondike goldfields, Yukon Territory, Canada: *Quaternary Research*, v. 66, p. 147–157, doi: 10.1016/j.yqres.2006.02.008.
- Schirmermeister, L., Kunitsky, V.V., Grosse, G., Kuznetsova, T.V., Derevyagin, A.Yu., Wetterich, S., and Siegert, C., 2008, The Yedoma Suite of the northeastern Siberian shelf region—Characteristics and concept of formation, in Kane, D.L., and Hinkel, K.M., eds., *Proceedings of the 9th International Conference on Permafrost*: University of Alaska Fairbanks, Institute of Northern Engineering, p. 1595–1601.
- Shapiro, B., Drummond, A.J., Rambaut, A., Wilson, M.C., Matheus, P.E., Sher, A.V., Pybus, O.G., Gilbert, M.T.P., Barnes, I., Binladen, J., Willerslev, E., Hansen, A.J., Baryshnikov, G.F., Burns, J.A., Davydov, S., Driver, J.C., Froese, D.G., Harington, C.R., Keddie, G., Kosintsev, P., Kunz, M.L., Martin, L.D., Stephenson, R.O., Storer, J., Tedford, R., Zimov, S., and Cooper, A., 2004, Rise and fall of the Beringian steppe bison: *Science*, v. 306, p. 1561–1565, doi: 10.1126/science.1101074.
- Sher, A.V., 1997, A brief overview of the Late Cenozoic history of the western Beringian lowlands, in Edwards, M.E., Sher, A.V., and Guthrie, R.D., eds., *Terrestrial paleoenvironmental studies in Beringia*: Fairbanks, Alaska Quaternary Center, p. 3–7.
- Sher, A.V., 1999, Traffic lights at the Beringian crossroads: *Nature*, v. 397, p. 103–104, doi: 10.1038/16341.
- Sher, A.V., Kuzimna, S.A., Kuznetsova, T.V., and Sulerzhitsky, L.D., 2005, New insights into the Weichselian environment and climate of the East Siberian Arctic, derived from fossil insects, plants, and mammals: *Quaternary Science Reviews*, v. 24, p. 533–569, doi: 10.1016/j.quascirev.2004.09.007.
- Sun, S.-s., and McDonough, W.F., 1989, Chemical and isotopic systematics of oceanic basalts: implications for mantle compositions and processes, in Saunders, A.D., and Norry, M.J., eds., *Magmatism in the ocean basins*: Geological Society of London Special Paper 42, p. 313–345.
- Weinstock, J., Willerslev, E., Sher, A., Tong, W., Ho, S.Y.W., Rubenstein, D., Storer, J., Burns, J., Martin, L., Bravi, C., Prieto, A., Froese, D., Scott, E., Xulong, L., and Cooper, A., 2005, Evolution, systematics, and phylogeography of Pleistocene horses in the New World: A molecular perspective: *PLoS Biology*, v. 3, p. e241, doi: 10.1371/journal.pbio.0030241.
- Westgate, J.A., Preece, S.J., Froese, D.G., Walter, R.C., Sandhu, A.S., and Schweger, C.E., 2001, Dating early and middle (Reid) Pleistocene glaciations in central Yukon by tephrochronology: *Quaternary Research*, v. 56, p. 335–348, doi: 10.1006/qres.2001.2274.
- Westgate, J.A., Preece, S.J., Froese, D.G., Pearce, N.J.G., Roberts, R.G., Demuro, M., Hart, W.K., and Perkins, W., 2008, Changing ideas on the identity and stratigraphic significance of the Sheep Creek tephra beds in Alaska and the Yukon Territory, northwestern North America: *Quaternary International*, v. 178, p. 183–209, doi: 10.1016/j.quaint.2007.03.009.

- White, J.M., Ager, T.A., Adam, D.P., Leopold, E.B., Liu, G., Jetté, H., and Schweger, C.E., 1997, An 18-million-year record of vegetation and climate change in northwestern Canada and Alaska; tectonic and global climatic correlates: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 130, p. 293–306, doi: 10.1016/S0031-0182(96)00146-0.
- Zazula, G.D., Froese, D.G., Schweger, C.E., Mathewes, R.W., Beaudoin, A.B., Telka, A.M., Harington, C.R., *an ternary Research*, v. 63, p. 189–198, doi: 10.1016/j.yqres.2004.11.003.
- Zazula, G.D., Froese, D.G., Elias, S.A., Kuzmina, S., La Farge, C., Reyes, A.V., Sanborn, P.T., Schweger, C.E., Smith, C.A.S., and Mathewes, R.W., 2006, Vegetation buried under Dawson tephra (25,300 ¹⁴C years B.P.) and locally diverse late Pleistocene paleoenvironments of Goldbottom Creek, Yukon, Canada: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 242, p. 253–286, doi: 10.1016/j.palaeo.2006.06.005.
- Zazula, G.D., Froese, D.G., Elias, S.A., Kuzmina, S., and Mathewes, R.W., 2007, Arctic ground squirrels of the mammoth-steppe: paleoecology of Late Pleistocene middens (~24,000–29,450 ¹⁴C yr BP), Yukon Territory, Canada: *Quaternary Science Reviews*, v. 26, p. 979–1003, doi: 10.1016/j.quascirev.2006.12.006.

Manuscript received 15 April 2009; accepted 27 May 2009. 